Primary Production in the Delta: Nutrient Considerations to Guide Conservation



Alexander E. Parker, PhD

California Maritime Academy

Many contributors to data collection and interpretation:

Dugdale / Wilkerson Lab Sacramento River Study

Richard Dugdale

Frances Wilkerson

Sarah Blaser

Victoria Hogue

Al Marchi

Adam Pimenta

Erica Kress

Jamie Lee

Allison Johnson

Chris Foe

Nutrient Ratios Project

Patricia Glibert

Jeff Alexander

Microcystis Project

Hans Paerl

Tim Otten

IRWM / Suisun Marsh Studies

Risa Cohen

Suisun Bay/ SWAMP

Karen Taberski


Primary Production in the Delta: Nutrient Considerations to Guide Conservation

The Delta is a mosaic of nutrient environments.

Experiments suggests that nutrient form, ratio and concentrations can alter algal community composition, nutrient and C uptake, and microbial function.

The relative importance of nutrient versus other drivers of patterns in primary production is likely site-specific. Modification of the Delta should consider potential nutrient-driven outcomes.

Thinking About Nutrients: The Ocean Perspective



From: Sigman and Hain 2012

Thinking About Nutrients: The Ocean Perspective

The availability of nitrogen limits primary production.

The distinction between "New" (i.e. nitrate-based) and "Regenerated" (ammonium-based) production is important in the supply and fate of primary production.

From: Sigman and Hain 2012

Thinking About Nutrients: The Ocean Perspective

Diatom-based food web based on "NEW" production.

Microbial loop favored through "REGENERATED" Production.

How do phytoplankton respond when nutrients are no longer limiting?

From: Sigman and Hain 2012

Phytoplankton Response to Anthropogenic Nutrients




200-L mesocosms amended with either NH₄ or NO₃ (50uM) and incubated for 72 hours.

>2-fold increase in chlorophyll-a, increase in DOC and bacterial activity

Phytoplankton Response to Anthropogenic

Nutrients







>2-fold increase in chlorophyll-a, increase in DOC and bacterial activity


Phytoplankton Response to Anthropogenic **Nutrients**

Increasing contribution by NH₄

Increasing N : P







Nutrients



smelt/bass/ silverside



phosphate



ammonium

Algae



diatoms



cryptophytes/flagellates



cvanobacteria

Copepods







Smelt/Silverside/Bass




Delta/longfin smelt




Young of Year striped bass




Glibert, 2010



Ephemeral spring and more persistent fall phytoplankton blooms in recent years.



2000 bloom associated with an improved light field, low ammonium concentrations and rapid nitrate uptake.,



Dugdale et al 2012


2010 diatom (*Entomoneis, Cyclotella, Melosira*) blooms lasted several weeks,

Phytoplankton increase 10-fold over 2009.



Zooplankton increase 9-fold over 2009

Delta smelt (70%) and longfin smelt (194%) increased (FMWT survey)





What Do We Know About Nutrients in the SFE?



Long history of measurements for the San Francisco Estuary (e.g. USGS).

Main stem of Sacramento and San Joaquin Rivers.


Paradigm is that nutrients are replete / not limiting.

Primary Production in the Delta: Nutrient Considerations to Guide Conservation

The Delta is a mosaic of nutrient environments.


Experiments suggests that nutrient form, ratio and concentrations can alter algal community composition, nutrient and C uptake, and microbial function.


The relative importance of nutrient versus other drivers of patterns in primary production is likely site-specific. Modification of the Delta should consider potential nutrient-driven outcomes.



From Parker & Cohen 2010


Large (4-fold) spatial and temporal variation in nutrients in Suisun Marsh.






From Parker & Cohen 2010


Large (4-fold) spatial and temporal variation in nutrients in Suisun Marsh.






From Parker & Cohen 2010


Large (4-fold) spatial and temporal variation in nutrients in Suisun Marsh.






From Parker & Cohen 2010

Large (4-fold) spatial and temporal variation in nutrients in Suisun Marsh.








From Parker & Cohen 2010

Large (>10-fold) variation over relatively small spatial scales.

Tidal wetlands in the confluence









From Parker et al. 2012





Summer N-limited in upper river and >4 fold variation in NH₄ seasonally.

Sacramento River



>4 fold variation in NH₄ seasonally.

Sacramento River

From SRWWTP, in Parker et al. 2010


	Station	Temp. C	Secchi (m)	NO ₃ +NO ₂ (μmol L ⁻¹)	NH ₄ (μmol L ⁻¹)	Urea-N (μmol L ⁻¹)	PO ₄ (μmol L ⁻¹)
Large River	RIO	21.2	1.5	10.22	12.6	0.44	1.02
	ANT	21.3	0.9	12.12	1.05	0.32	1.05
Small River	МОК	21.6	1.9	9.36	3.61	0.64	0.43
	OLD	22.8	3.5	4.59	1.34	0.14	1.66
Flooded Island	MIL	23.1	2.3	21.64	0.32	0.56	1.70
	FRK	21.6	>2	10.40	1.15	0.78	1.56

Central and South Delta

The Delta is a mosaic of nutrient environments.


Experiments suggests that nutrient form, ratio and concentrations can alter algal community composition, nutrient and C uptake, and microbial function.

The relative importance of nutrient versus other drivers of patterns in primary production is likely site-specific. Modification of the Delta should consider potential nutrient-driven outcomes.



Elevated chlorophyll-a associated with NO₃ uptake

From Dugdale et al. 2007



Ammonium (NH₄) inhibits phytoplankton access to nitrate (NO₃)

NH₄ \approx 4 μ M (0.056 mg/L) inhibits NO₃ uptake.

NH₄ \approx 1 μ M (0.014 mg/L) inhibits NO₃ uptake by 60%


From Dugdale et al. 2007

NH₄ inhibition of NO₃ uptake is not universal (nor are threshold NH₄ concentrations) across phytoplankton taxa (Dortch 1990).

NH₄ inhibits transport of NO₃ into the cell.

ln(Y) = -1.28 * ln(X) - 4.26

NH₄ inhibits nitrate reductase (a necessary step in NO₃ assimilation into protein) activity and specificity.




From Dugdale et al. 2007


0.03


Nutrients and Primary Production

in the Delta











Nutrients and Primary Production

in the Delta


Chlorophyll-a is dominated by diatoms









Nutrients and Primary Production in the Delta – *Microcystis*




Nutrients and Primary Production in the Delta – *Microcystis*






Nutrients and Primary Production in the Delta – *Microcystis*



Parker et al in nren




Heil, Revilla, Glibert, Murasko

Sacramento Gareta Bend WTP Sacramento Regional Elk Grove Vacaville Kenady Landing Galt Vista * Isleton Confluence Qakley Light 14 Antioch Stockton Brentwood Stockton WTP Marsh Creek Discovery 8 ALGAL CELLS Brandt Bridge millions/L Manteca Centric diatoms Pennate diatoms Cryptophytes Chlorophytes Blue-green algae Flagellates 15 Miles

From Pulse of the Estuary 2012; Based on Kress 2012

Nutrients and Primary Production

in the Delta



From Pulse of the Estuary 2012; Based on Kress 2012

The Challenge: Placing Experimental Results in an Ecological Framework

The Delta is a mosaic of nutrient environments.

Experiments suggests that nutrient form, ratio and concentrations can alter algal community composition, nutrient and C uptake, and microbial function.

The relative importance of nutrient versus other drivers of patterns in primary production is likely site-specific. Modification of the Delta should consider potential nutrient-driven outcomes.

Primary Production in the Delta: Nutrient Considerations to Guide Conservation

Funding from:

The Delta Science Program

Water Quality Control Boards (R2, R5)

The State and Federal Contractors Water Agency

CA Dept. Fish and Wildlife

Thanks to:

Sacramento Regional County Sanitation District

R/V Questuary










Delta Stewardship Council






The Challenge: Placing Experimental Results in an Ecological Framework


Ecological Modelling 263 (2013) 291-307



Contents lists available at SciVerse ScienceDirect

Ecological Modelling





A biogeochemical model of phytoplankton productivity in an urban estuary: The importance of ammonium and freshwater flow



Richard C. Dugdale*, Frances P. Wilkerson, Alexander E. Parker

Romberg Tiburon Center, San Francisco State University, 3152 Paradise Drive, Tiburon, CA 94920, United States

